

# The bryophytes of the spruce forests of South Tyrol: species list, distribution and ecology

## Abstract

The knowledge about the bryophytes of forest habitats in the Alps, and more particularly in South Tyrol, is still inadequate. Most works are floristic, and scarce are quantitative studies examining spruce forests. Therefore, aims of this work are: 1) to increase the knowledge of bryophyte distribution in the spruce forests of South Tyrol; 2) to discuss the ecology and distribution of the most significant species or species groups. A total of 48 plots were surveyed in South Tyrol along the elevational gradient and covering the province in 8 main areas. Bryophyte species were examined on three different types of substrates, tree trunks of spruce (*Picea abies*), deadwood and on the forest floor. This work allowed to record 120 species (91 mosses and 29 liverworts), 27 considered rare or threatened. One interesting discovery was *Sciuro-hypnum curtum*, a pleurocarpous moss firstly recorded in Italy. The ecology of the most significant groups of species were outlined. Elevation, canopy closure, rainfall and amount of necromass were the main factors determining the species distribution. The role of the environment and the responsibility of forest managers for the biodiversity conservation was discussed.

Keywords: bryophytes, environmental gradients, liverworts, mosses, South Tyrol

## Introduction

The concept of “sustainable forest management” considers the maintenance of biodiversity a crucial point, and it is a target of many governments, scientific and economic initiatives (LINDENMAYER et al. 2000). The sustainable forest management is included in the more encompassing ecological target of “forest integrity”. Forest integrity is the capacity to support and maintain a balanced, integrated, adaptive community of organisms having a species composition, diversity and functional organization comparable to that of similar undisturbed forest ecosystem (e.g. TIERNEY et al. 2009). However, defining criteria for forest integrity is challenging.

Bryophytes are important components of forest integrity. Their contribute to species richness is at least as high as that of the vascular plants in many forest types (e.g. GRYTNES et al. 2006). Although the live biomass of bryophytes comprises a small fraction of the total ecosystem carbon content, they greatly influences the function of the ecosystem (OECHEL & VAN CLEVE 1986, KOLARI et al. 2006). Bryophytes are involved in the processes regulating soil thermal regimes, hydrology and nutrient availability (SVEINBJORNSSON & OECHEL 1992).

The relationship between bryophyte species occurrence and habitats in the spruce forests is well studied. Some investigations focussed on the occurrence of species on selected microhabitats (SÖDERSTRÖM 1988, MÁRIALIGETI et al. 2009, KIRÁLY & ÓDOR 2010), others examined the variation in bryophyte community along environmental gradients and scales (MILLS & MACDONALD 2005, TINYA et al. 2009). However, even though studies

## Author's address

Daniel Spitale  
Museo di Scienze Naturali  
dell'Alto Adige,  
via Bottai 1  
39100 Bolzano  
spitale.daniel@gmail.com

submitted: 30. 04. 2015  
accepted: 01. 10. 2015

dealing with the relationships between bryophytes and environmental and management factors are not rare, more patchy are the regional distribution of such works. This is an important shortcoming, as bioindicators often are specific of a particular pool of species in a biogeographic area.

The knowledge about the bryophytes of spruce forests in the Alps, and more particularly in South Tyrol, is scarce. Most works are floristic and considered the spruce forests only among other habitats (e.g. ALEFFI & CORTINI PEDROTTI 1996). In South Tyrol bryological studies date back to the 19<sup>th</sup> century, the most important work represented by DALLA TORRE & SARNTHEIN (1904). Later, almost all the knowledge about the bryophyte distribution in the province has been collected by Ruprecht Düll, and summarized in the check-list of South Tyrol (DÜLL 2006). However, even though the province seems to be well studied from the floristic point of view (in contrast to other Italian provinces), still lacking are quantitative studies examining spruce forests. Therefore, the aims of this work are: 1) increasing the knowledge of bryophyte distribution in the spruce forests of South Tyrol; 2) discuss the ecology and distribution of the most significant species or species groups. This contribution will enhance our comprehension of the bryodiversity and will elucidate the peculiarities of this taxonomic group in South Tyrol.

## Methods

Field sampling was preceded by an accurate examination of available data in collaboration with the Department of Forest Planning of Bolzano. Spruce-forests are the most common type of wood in the province covering a total of 3330 km<sup>2</sup> (AA. VV. 2010). They represent the 88% of the total forested area. The first localization of the potential survey areas was accomplished balancing the number of plots according to the distribution in the province. Eight continuous gradients from 900 m to 1900 m a.s.l. were selected (Figure 1). The gradient was distinguished into three elevational steps (900-1200 m a.s.l. = sub-montane, 1400-1600 m = montane and 1800-1900 m = sub-alpine). At each elevational steps, two circular plots (13 m radius) were allocated randomly once the following conditions were fulfilled: (i): the property of the forest had to be public; (ii) the spruce trees had to be adult and ready to be harvested; (iii) the plots had to be distant hundreds of meters (mean distance between plots within elevational steps: 574 m). In each plot, bryophytes were sampled on three types of substrates: tree trunks of living spruces (*Picea abies*), deadwood and forest floor. We used standard grids of 10 x 50 cm to sample bryophytes. A “sample” consisted of 4 grids within which the presence of the species was recorded. This kind of grid is commonly used also by lichenologists (NASCIMBENE 2014). On tree trunks, the grids were placed at North and South both on the trunk base and at 1 m height (4 grids = 200 cm<sup>2</sup>). Deadwood (stumps and fallen logs), were selected only if the surface area exceeded 4 grids (50 cm<sup>2</sup> x 4 = 200 cm<sup>2</sup>). Because most of the deadwood consisted of stumps, two grids were placed on the vertical side and two on the top. Ground substrate was similarly sampled arranging 4 grids side by side and recording the presence of the species. A total of 5 samples for each substrate were surveyed on each plot. Sampling work was completed in the 2013.

Canopy closure (the proportion of sky hemisphere obscured by vegetation), influences the forest floor microclimate and light conditions (KORHONEN et al. 2006). To estimate canopy closure we used a spherical densiometer (it is a convex mirror, engraved with 24 squares). The densiometer was used by holding it at breast height and counting the number of squares occupied by the canopy reflected in the convex mirror (KORHONEN et al. 2006). Canopy closure was measured above each bryophyte sample for a total of 15 points per plot. The average of them represented the canopy closure of the plot.

The mean age of the trees on the plot was quantified by extracting cores using a Pressler-type increment borer at a vertical height of 1.30 m (NASCIMBENE et al. 2014). The procedure was repeated on the 5 trees where corticolous species were surveyed.




Figure 1. Distribution of plots surveyed between 2013-2014 in South Tyrol, Italy. The eight areas are identified with the following letters: P = Prato allo Stelvio / Prad am Stilfserjoch, Si = Silandro / Schlanders, L = Laces / Latsch; T = Tires / Tiers, F = Funes / Villnöß, M = Monguelfo / Welsberg; Se = Selva dei Molini / Mühlwald; R = Racines / Ratschings. Symbols represent plots.

The amount of deadwood in the plots was estimated inventorying all the stumps and downed logs with dbh > 7 cm. The volume of deadwood was calculated according to the Huber formula (HUSH et al. 2003). The decay stage of stumps was classified according to a four-stage system adopted by MOTTA et al. (2006): (1) bark intact, wood hard; (2) bark almost completely intact, wood hard in the outermost part of stump, decaying in the innermost part, texture with large pieces; (3) only traces of bark, decay spread in most of the stump, texture with blocky pieces; (4) bark absent, wood soft and with powdery structure. The decay stage of logs was determined using again four classes: (1) bark intact, shape round, wood texture intact; (2) trace of bark, shape round and wood still hard; (3) bark absent, shape round to oval, texture of wood with blocky pieces; (4) no bark, shape oval, wood soft and powdery structure.

Mosses were identified mainly according to the keys of CORTINI PEDROTTI (2001, 2006), SMITH (2004) and NYHOLM (1986), while PATON (1999) and DAMSHOLT (2002) were used for liverworts.

The data analyses performed were as follows: Species diversity was calculated both as number of species and also using Shannon Index. In order to get the habitat preference of species, weighted average and tolerance was calculated for several environmental variables. Canonical Correspondence Analysis was used to identify the main environmental factors driving the species assemblages. All the analyses were accomplished with the R statistical platform (R DEVELOPMENT CORE TEAM 2014).

## Results and discussion

Overall, 120 species of mosses and liverworts were found in 48 plots of spruce forests. On average, there were 20 species on each plot (range 9-37), among which 6 liverworts (range 0-15 species). Shannon diversity was on average 2.6 (0.32 SD). The number of bryophytes recorded on three types of substrates (on tree trunks of spruces, on deadwood and on the forest floor) was very similar to the number of lichens recorded only on trees of the same plots (120 vs 124, NASCIMBENE & MARINI 2015). Comparison of these values with other studies which used different sampling strategies cannot be done safely.

A total of 26 species were included in the Red List of the neighbor Province of Trento (CORTINI PEDROTTI & ALEFFI 2011), the most significant species were: *Calypogeia neesiana* (EN), *Cynodontium gracilescens* (VU), *Cynodontium strumiferum* (VU), *Lepidozia reptans* (VU), *Lophozia ascendens* (VU), *Lophozia longiflora* (EN), *Lophozia ventricosa* (VU), *Plagiothecium denticulatum* (VU), *Plagiothecium succulentum* (VU), *Pohlia longicollis* (VU). According to DÜLL (2006) 17 species are considered rare, and 10 extremely rare or recorded few times in the province. An interesting discovery was *Sciuro-hypnum curtum*, a pleurocarpous moss firstly recorded in Italy (SPITALE 2015a). The species has been synonymized with *S. oedipodium* but IGNATOV & MILYUTINA (2007) showed that they are two distinct species. As currently understood, *S. curtum* is a widespread species in boreal forests of Eurasia and occurs also in the east of North America. Instead, *Sciuro-hypnum oedipodium* is mainly an American species, with few localities in Eurasia (Caucasus and in Chukotka). Another interesting finding, already known in the province but only with scanty data, was *Buxbaumia viridis*. To this species we devoted a special study to clarify its ecology and distribution (SPITALE & MAIR 2015, SPITALE et al. 2015).

### Liverworts

Among the most characteristic taxa of liverworts found in the spruce forests of South Tyrol were *Barbilophozia* (*B. barbata*, *B. hatcheri* and *B. lycopodioides*), *Lophozia* (*L. ascendens*, *L. incisa*, *L. longidens*, *L. longiflora*, *L. silvicola*, *L. ventricosa*), *Calypogeia* (*C. integristipula*, *C. muelleriana*, *C. neesiana*, *C. suecica*) and *Tritomaria* (*T. exsecta*, *T. exsectiformis*).

The species of *Barbilophozia* were quite common but they were characteristic of intermediate elevation (1300-1600 m a.s.l.). With the exception of *B. lycopodioides* which prefers the forest floor, the other two species inhabit tree trunks, deadwood and forest floor indifferently.

The genus *Lophozia* is a difficult taxon with many species with overlapped morphological characters (DAMSHOLT 2002). Often they have abundant gemmae, green in *L. ventricosa*, *L. incisa*, *L. ascendens*, *L. silvicola* and red in *L. longidens*. *Lophozia incisa* and *L. longiflora* were found only on deadwood, *L. longidens*, *L. silvicola*, *L. ventricosa* both on tree trunks and deadwood. All the *Lophozia* species preferentially grow between 1500 and 1800 m a.s.l.

Also the genus *Calypogeia* is a difficult taxon which needs abundant and well developed specimens to be determined at species level with some confidence (BUCZKOWSKA 2004). In particular, the couple *C. integristipula*-*C. neesiana* (underleaves entire) and *C. muelleriana* - *C. suecica* (underleaves with distinct sinus) can be sometimes difficult to separate. The *Calypogeia* species occur preferentially between 1650 m and 1850 m a.s.l., and on forests with abundant deadwood on advanced stage of decay (class 3 and 4).

The genus *Tritomaria*, with abundant and red gemmae occurring at the top of the shoots, is readily recognizable in the field. However, in order to separate the two species, gemmae should be checked under the microscope to inspect their shape. *Tritomaria exsectiformis* was found only two times, whereas *T. exsecta* was found in 6 plots. They prefer elevation between 1500 and 1700 m a.s.l. and they grow on deadwood.



## Mosses

The taxon *Hypnum cupressiforme* probably is among the few, most emblematic species of forests (Fig. 2). It is a polymorphic species, often treated as a species complex. It can grow anywhere in the spruce forests, and often it was the only species able to colonize the acid bark of spruces. It occupied a well-defined range of altitude with an optimum at 1274 m a.s.l (tolerance 254 m). *Hypnum andoi* shared the same habitat condition of *H. cupressiforme* and it is not so rare as previously retained (DÜLL 2006). *Hypnum pallens* is a rare species (also according to DÜLL 2006), and in fact it was found only once.

The genus *Plagiothecium* was well represented with five species but only two were common (*P. denticulatum* and *P. laetum*). *Plagiothecium nemorale*, *P. platyphyllum* and *P. succulentum* are rare species (one record each) even though DÜLL (2006) stated that they are sporadic. Probably, the optimal habitat is not the spruce forest in South Tyrol. Concerning the altitudinal gradient occupied by the two more common species, *P. denticulatum* has an optimum at 1521 m (tolerance 384 m), whereas *P. laetum* stays upward (optimum at 1707 m, tolerance 212 m). *Plagiothecium denticulatum* grows both on the forest floor and on deadwood, more rarely on tree trunks; *P. laetum* occurs indifferently on all the substrates.

The genus *Polytrichum* s.l. (including also *Polytrichastrum*), is another widespread taxon but with several species differentiation. For example, *P. formosum* and *P. alpinum* were the most common, but occupied different altitudinal ranges (lower the former, upper the latter). Both the species have been never found on tree trunks. *Polytrichum longisetum* was retained extremely rare by DÜLL (2006), and in fact it has been found only in 3 plots.

The genus *Dicranum* was dominated by two species (*D. montanum* and *D. scoparium*) which occupy indifferently all the substrates and altitudes. The other two species, *D. brevifolium* and *D. muehlenbeckii* were singletons and are not typical for forest habitats. Identification of *D. brevifolium* and *D. muelenbeckii* should be considered tentative, as intermediate forms exists (HEDENÄS & BISANG 2004).



Figure 2. The moss *Hypnum cupressiforme*, a common species colonizing all the substrates in forests with an optimum growth and diffusion around the 1300 m s.l.m.

## General patterns

In order to obtain a general overview of the main environmental variables driving the bryophyte distribution in the spruce forests, a Canonical Correspondence Analysis (CCA) was performed (Fig. 3). The analysis suggested that the most important factors were elevation, age, canopy closure, rainfall and the amount of necromass in advanced stage of decay (class 3 and 4). The model was highly significant ( $F = 2.53$ ,  $P < 0.001$ ). Elevation and age (average of spruce trees in the surveyed plot) had similar sign and importance in determining species distribution, and both factors were opposite to canopy closure. These three environmental variables explained the turnover of species along the altitudinal gradient. Tree age co-varied with elevation as a result of different tree growth rates along the elevation gradient. That is, trees with similar diameter were on average older at high elevation. Canopy closure was negatively correlated with elevation and age, because mature forests and upland forests tend to have more gaps. Thus, the primary factors determining the bryophyte species composition were elevation (and co-related factors like temperature), and light (SPITALE 2015b). These both factors gain importance as the elevation gradient extends. These fundamental environmental variables are directly related to the physiology of bryophytes. In lowland forests, which have higher temperatures, the thallus of bryophytes desiccate more rapidly than upland (they are poikilohydric organisms, PROCTOR 2000). This environmental setting probably is too stressful for many species, and species diversity decreases in lowland forests. Only species more tolerant and with large colony like *Hypnum cupressiforme*, *Isoetecium alopecuroides*, *Dicranum scoparium*, are able to dominate at low elevation. A comparable pattern was observed also on lichens for the same study sites (NASCIMBENE & MARINI 2015). Rainfall was the third most important environmental factor according to the CCA analysis (Fig. 3). It was orthogonal to elevation, so it accounted for another fraction of the total variance. In ecological terms, this suggested some spatial differences in the assemblage composition related to the different rainfall observed in South Tyrol (563–1182 mm/yr). Even rainfall was related to fundamental physiological processes of bryophytes. Finally, the amount of necromass was able to explain a significant amount of variance. Necromass constitute a special habitat, crucial for many organisms in the forests other than bryophytes (JONSSON et al. 2005). A number of liverworts and saproxylic species like *Buxbaumia viridis* (SPITALE & MAIR 2015) were directly dependent from the amount of deadwood in forests. This result alone suggested the fundamental role that forest managers play in determining the sustainable use of the forests. Unfortunately, based on the data we collected, it seems that on average the amount of deadwood was largely insufficient to sustain the forest integrity (SPITALE et al. 2015). On average, 15 m<sup>3</sup>/ha (range 1–53 m<sup>3</sup>/ha) of deadwood was present in the spruce forests of South Tyrol. As suggested also by MÜLLER & BÜTLER (2010), this amount is below the threshold suggested to be effective in maintaining the integrity of spruce forests (20–30 m<sup>3</sup>/ha).

Figure 3. Canonical Correlation Analysis on the bryophytes of spruce forests of South Tyrol. Canopyclos = canopy closure, necrom.cl3.4 = necromass in decay class 3 and 4.




Table 1. List of the species recorded in the 48 spruce forests in South Tyrol in 2013. Rec = number of records; s, w, t, are the relative frequency of the species respectively on the soil, deadwood and tree trunks. RL = Red List species compiled according to Cortini Pedrotti & Aleffi (2011) concerning the neighbor province of Trento; distr = species distribution according to Düll (2006): v = verbreitet = common; h = häufig = frequent; zv = ziemlich verbreitet = fairly widespread z-v = meint stellenweise = sporadic but locally common; z = zerstreut = sporadic; r-z = selten bis zerstreut = from sporadic to rare; r = selten = rare; rr = nur an einem oder sehr wenigen Orten nachgewiesen = recorded only once or in few places. Areas were referred to the location of surveyed plots: P = Prato allo Stelvio; Si = Silandro; L = Laces, T = Tires, F = Funes, M = Monguelfo, Se = Selva del Molini; R = Racines. Altitude was the weighted average. Nomenclature was according to Ros et al. (2007, 2013).

	REC	S	W	T	RL	DISTR	AREAS	ALTITUDE
<i>Abietinella abietina</i> (Hedw.) M.Fleisch.	3	0.9	0.1	0.0		z-v	Si, L	1004
<i>Alleniella complanata</i> (Hedw.) S. Olsson, Enroth & D. Quandt	1	1.0	0.0	0.0		z-v	P	1110
<i>Antitrichia curtipendula</i> (Hedw.) Brid.	3	1.0	0.0	0.0		z	Si, L	1057
<i>Atrichum undulatum</i> (Hedw.) P.Beauv.	1	1.0	0.0	0.0		v	M	1880
<i>Barbilophozia barbata</i> (Schmidel ex Schreb.) Loeske	8	0.5	0.3	0.2	NT	z	P, L, T, Se, F	1334
<i>Barbilophozia hatcheri</i> (A. Evans) Loeske	15	0.4	0.3	0.3		r	P, Si, L, M, Se, R, F	1556
<i>Barbilophozia lycopodioides</i> (Wallr.) Loeske	11	0.8	0.1	0.1	NT	z	P, M, Se, R, F	1637
<i>Bartramia ithyphylla</i> Brid.	2	1.0	0.0	0.0		z	Si, M	1540
<i>Bazzania trilobata</i> (L.) Gray	1	0.0	1.0	0.0		z	R	1470
<i>Blepharostoma trichophyllum</i> (L.) Dumort.	25	0.3	0.6	0.1	NT	z	P, Si, L, T, M, Se, R, F	1630
<i>Brachytheciastrum velutinum</i> (Hedw.) Ignatov & Huttunen	34	0.5	0.3	0.2		v	P, Si, L, T, M, Se, R, F	1458
<i>Brachythecium rutabulum</i> (Hedw.) Schimp.	2	0.4	0.6	0.0		v	P, F	1120
<i>Brachythecium salebrosum</i> (Hoffm. ex F.Weber & D.Mohr) Schimp.	11	0.6	0.4	0.0		z	Si, L, M, Se, R, F	1752
<i>Bryoerythrophyllum recurvirostrum</i> (Hedw.) P.C.Chen	1	0.0	1.0	0.0		z-v	T	1803
<i>Bryum elegans</i> Nees	3	0.5	0.0	0.5		z	M, F	1679
<i>Buxbaumia viridis</i> (Moug. ex Lam. & DC.) Brid. ex Moug. & Nestl.	3	0.3	0.7	0.0		z	M, Se, F	1422
<i>Calypogeia integristipula</i> Steph.	10	0.2	0.8	0.0		r	P, L, M, Se, R, F	1800
<i>Calypogeia muelleriana</i> (Schiffn.) Müll. Frib.	1	0.0	1.0	0.0		z	R	1800
<i>Calypogeia neesiana</i> (C. Massal. & Carestia) Müll. Frib.	9	0.3	0.7	0.0	EN	r	L, M, Se, R, F	1673
<i>Calypogeia suecica</i> (Arnell & J. Perss.) Müll. Frib.	3	0.8	0.2	0.0		rr	M, Se, F	1847
<i>Campylium protensum</i> (Brid.) Kindb.	2	1.0	0.0	0.0		z	M, F	1853
<i>Campylophyllum halleri</i> (Hedw.) M.Fleisch.	2	1.0	0.0	0.0		z	F	1853
<i>Campylopus pyriformis</i> (Schultz) Brid.	1	1.0	0.0	0.0	VU	rr	T	1188
<i>Cephalozia bicuspidata</i> (L.) Dumort.	3	0.5	0.5	0.0		z	Se, R, F	1736
<i>Ceratodon purpureus</i> Brid.	4	0.3	0.5	0.3		h	P, Si, L	1459
<i>Cirriphyllum piliferum</i> (Hedw.) Grout	2	1.0	0.0	0.0		z	M, F	1410
<i>Ctenidium molluscum</i> (Hedw.) Mitt.	2	1.0	0.0	0.0		v	T, F	1630
<i>Cynodontium gracilescens</i> (F.Weber & D.Mohr) Schimp.	5	0.2	0.6	0.2	VU	r	Si, L, M, Se	1602
<i>Cynodontium polycarpon</i> (Hedw.) Schimp.	5	0.4	0.4	0.2		z	P, Si, L	1532
<i>Cynodontium strumiferum</i> (Hedw.) Lindb.	4	0.1	0.7	0.1	VU	z	P, Se	1549
<i>Dicranodontium denudatum</i> (Brid.) E.Britton	2	0.3	0.7	0.0	NT	z	T, R	1627
<i>Dicranum brevifolium</i> (Lindb.) Lindb.	1	0.0	1.0	0.0		rr	L	1900
<i>Dicranum montanum</i> Hedw.	40	0.1	0.5	0.4	NT	z	P, Si, L, T, M, Se, R, F	1582
<i>Dicranum muehlenbeckii</i> Bruch & Schimp.	1	0.0	1.0	0.0		z-v	L	1820
<i>Dicranum scoparium</i> Hedw.	48	0.3	0.4	0.2		z-v	P, Si, L, T, M, Se, R, F	1586
<i>Encalypta ciliata</i> Hedw.	1	1.0	0.0	0.0		sz	Si	1200
<i>Eurhynchium angustirete</i> T.J.Kop.	8	0.8	0.2	0.1	EN	v	T, M, R, F	1147
<i>Eurhynchium pulchellum praecox</i> (Hedw.) Ochyra & Żarnowiec	1	1.0	0.0	0.0		z	Si	1200
<i>Eurhynchium striatum</i> (Hedw.) Schimp.	1	1.0	0.0	0.0		r	T	1188
<i>Exsertotheca crispa</i> (Hedw.) S. Olsson, Enroth & D. Quandt	1	1.0	0.0	0.0		z	T	1188

	REC	S	W	T	RL	DISTR	AREAS	ALTITUDE
<i>Fissidens adianthoides</i> Hedw.	4	0.6	0.4	0.0		r	T, F	1347
<i>Fissidens taxifolius</i> Hedw.	1	1.0	0.0	0.0		z	M	1880
<i>Fissidens viridulus</i> (Sw. ex anon.) Wahlenb.	1	1.0	0.0	0.0		rr	Si	1180
<i>Frullania dilatata</i> (L.) Dumort.	11	0.1	0.2	0.8		v	P, Si, L, T, R	1235
<i>Hedwigia ciliata</i> (Hedw.) P.Beauv.	3	0.4	0.4	0.2		v	L	1088
<i>Herzogiella seligeri</i> (Brid.) Z.Iwats.	21	0.1	0.7	0.1		z	P, Si, L, T, M, Se, R, F	1445
<i>Heterocladium dimorphum</i> (Brid.) Schimp.	3	0.5	0.3	0.2		r-z	M, R, F	1646
<i>Homalothecium lutescens</i> (Hedw.) H.Rob.	1	0.0	1.0	0.0		z-v	T	1540
<i>Hylocomium splendens</i> (Hedw.) Schimp.	48	0.7	0.3	0.0		z-v	P, Si, L, T, M, Se, R, F	1449
<i>Hymenoloma crispulum</i> (Hedw.) Ochyra	2	1.0	0.0	0.0		z-v	P, Se	1818
<i>Hypnum andoi</i> A.J.E.Sm.	13	0.0	0.4	0.6		rr	Si, L, T, M, Se, F	1275
<i>Hypnum cupressiforme</i> Hedw.	37	0.2	0.4	0.4		v-h	P, Si, L, T, M, Se, R, F	1279
<i>Hypnum pallescens</i> (Hedw.) P.Beauv.	1	0.0	0.0	1.0		rr	P	1820
<i>Isothecium alopecuroides</i> (Lam. ex Dubois) Isov.	12	0.5	0.2	0.3		z-v	P, Si, L, T, Se, F	1316
<i>Lejeunea cavifolia</i> (Ehrh.) Lindb.	1	1.0	0.0	0.0		z	Se	1200
<i>Lepidozia reptans</i> (L.) Dumort.	22	0.2	0.7	0.1		z	P, Si, T, M, Se, R, F	1596
<i>Lophocolea heterophylla</i> (Schrad.) Dumort.	32	0.3	0.3	0.4		z	P, Si, L, T, M, Se, R, F	1622
<i>Lophozia ascendens</i> (Warnst.) R. M. Schust.	1	0.0	1.0	0.0	VU	rr	Se	1550
<i>Lophozia incisa</i> (Schrad.) Dumort.	8	0.0	1.0	0.0	NT	z	M, Se, F,	1734
<i>Lophozia longidens</i> (Lindb.) Macoun	13	0.0	0.4	0.6		r	P, Si, L, M, Se, F	1639
<i>Lophozia longiflora</i> (Nees) Schiffn.	7	0.2	0.8	0.0	EN	r	M, Se, R, F	1811
<i>Lophozia silvicola</i> H. Buch	5	0.3	0.5	0.2		z	P, M, Se, F	1588
<i>Lophozia ventricosa</i> (Dicks.) Dumort.	6	0.2	0.4	0.4	VU	r	P, Si, L, M, Se	1631
<i>Metzgeria furcata</i> (L.) Dumort.	3	0.8	0.0	0.2		v	P, Se, R	1145
<i>Mnium lycopodioides</i> Schwägr.	1	1.0	0.0	0.0		r	M	1880
<i>Mnium spinosum</i> (Voit) Schwägr.	24	0.8	0.2	0.1		z-v	P, Si, L, T, M, Se, R, F	1495
<i>Mnium stellare</i> Hedw.	2	1.0	0.0	0.0		z	L, M	1524
<i>Mnium thomsonii</i> Schimp.	2	0.5	0.0	0.5		z	Si, F	1695
<i>Orthotrichum pallens</i> Bruch ex Brid.	1	0.0	0.0	1.0		z	P	1110
<i>Orthotrichum scanicum</i> Gronvall	1	0.0	1.0	0.0	VU	rr	Si	1490
<i>Orthotrichum speciosum</i> Nees	2	0.0	0.5	0.5		z	P, Si	1460
<i>Orthotrichum striatum</i> Hedw.	2	0.0	1.0	0.0		z-v	Si L	1213
<i>Oxystegus tenuirostre</i> (Hook. & Taylor) A.J.E.Sm.	1	1.0	0.0	0.0	VU	z	M	1880
<i>Paraleucobryum longifolium</i> (Hedw.) Loeske	3	0.7	0.0	0.3	VU	z-v	L, Se	1214
<i>Pellia neesiana</i> (Gottsche) Limpr.	2	1.0	0.0	0.0	VU	r	M	1877
<i>Plagiochila asplenoides</i> (L. emend. Taylor) Dumort.	9	0.5	0.2	0.2		z	M, Se, F	1377
<i>Plagiochila porelloides</i> (Torrey ex Nees) Lindenb.	19	0.6	0.3	0.1		v	P, Si, T, M, R, F	1483
<i>Plagiomnium affine</i> (Blandow ex Funck) T.J.Kop.	8	0.6	0.3	0.1		z	T, M, Se, F	1196
<i>Plagiomnium ellipticum</i> (Brid.) T.J.Kop.	1	1.0	0.0	0.0	VU	r	M	1880
<i>Plagiomnium undulatum</i> (Hedw.) T.J.Kop.	5	0.7	0.2	0.1		v	P, M, F	1072
<i>Plagiothecium denticulatum</i> (Hedw.) Schimp.	10	0.6	0.3	0.1	VU	z	L, T, M, Se, R, F	1521
<i>Plagiothecium laetum</i> Schimp.	25	0.2	0.3	0.5		z	P, Si, L, T, M, Se, R, F	1707
<i>Plagiothecium nemorale</i> (Mitt.) A.Jaeger	1	1.0	0.0	0.0		z	M	1880
<i>Plagiothecium platyphyllum</i> Mönk.	1	0.0	1.0	0.0			F	950
<i>Plagiothecium succulentum</i> (Wilson) Lindb.	1	0.0	1.0	0.0	VU	z	F	950



	REC	S	W	T	RL	DISTR	AREAS	ALTITUDE
<i>Pleurozium schreberi</i> (Willd. ex Brid.) Mitt.	31	0.7	0.3	0.0		z-v	P, Si, L, T, M, Se, R, F	1612
<i>Pogonatum umigerum</i> (Hedw.) P.Beauv.	2	0.5	0.5	0.0		z-v	M, F	1385
<i>Pohlia cruda</i> (Hedw.) Lindb.	4	1.0	0.0	0.0		z	P, Si, M	1482
<i>Pohlia longicolla</i> (Hedw.) Lindb.	2	0.5	0.5	0.0	VU	r	T, R	1811
<i>Pohlia nutans</i> (Hedw.) Lindb.	16	0.3	0.7	0.0		z-v	P, Si, L, T, M, Se, F	1656
<i>Polytrichastrum alpinum</i> (Hedw.) G.L.Sm.	11	0.6	0.4	0.0		z-v	P, Si, L, M, Se, R	1851
<i>Polytrichum commune</i> Hedw.	1	0.7	0.3	0.0		z	Se	1815
<i>Polytrichum formosum</i> Hedw.	12	0.3	0.3	0.3		z-v	T, M, Se, R, F	1424
<i>Polytrichum juniperum</i> Hedw.	6	1.0	0.0	0.0		z-v	L, M, Se, R	1672
<i>Polytrichum longisetum</i> Sw. ex Brid.	3	0.6	0.4	0.0		rr	M, R, F	1742
<i>Polytrichum piliferum</i> Hedw.	2	0.5	0.5	0.0		v	P, M	1650
<i>Pseudoscleropodium purum</i> (Hedw.) M.Fleisch.	1	1.0	0.0	0.0		z-v	T	1188
<i>Pterigynandrum filiforme</i> Hedw.	14	0.3	0.2	0.5		z-v	P, Si, L, M, Se, R	1407
<i>Ptilidium pulcherrimum</i> (Weber) Vain.	17	0.0	0.4	0.5		z	P, T, M, Se, R, F	1541
<i>Ptilium crista-castrensis</i> (Hedw.) De Not.	2	0.5	0.5	0.0		z-v	Se, F	1213
<i>Ptychostomum pallens</i> (Sw.) J.R. Spence	12	0.5	0.3	0.2		z	P, Si, L, T, M, F	1470
<i>Racomitrium microcarpon</i> (Hedw.) Brid.	1	1.0	0.0	0.0	NT	r	Se	1870
<i>Radula complanata</i> (L.) Dumort.	12	0.3	0.2	0.5		v	P, Si, L, T, M, Se, R, F	1227
<i>Rhizomnium magnifolium</i> (Horik.) T.J.Kop.	1	1.0	0.0	0.0	CR	r	Se	1815
<i>Rhizomnium punctatum</i> (Hedw.) T.J.Kop.	9	0.4	0.5	0.1		z-v	M, Se, R, F	1501
<i>Rhytidiadelphus squarrosus</i> (Hedw.) Warnst.	2	1.0	0.0	0.0		sz	M, F	1070
<i>Rhytidiadelphus triquetrus</i> (Hedw.) Warnst.	44	0.8	0.2	0.0		v	P, Si, L, T, M, Se, R, F	1425
<i>Sanionia uncinata</i> (Hedw.) Loeske	18	0.2	0.7	0.1		z-v	P, Si, L, T, M, Se, R, F	1689
<i>Scapania aequiloba</i> (Schwägr.) Dumort.	2	1.0	0.0	0.0		r	F	1853
<i>Schistidium robustum</i> (Nees & Hornsch.) H.H.Blom	1	1.0	0.0	0.0		rr	F	1860
<i>Sciuro-hypnum curtum</i> (Lindb.) Ignatov	1	1.0	0.0	0.0		new	L	1820
<i>Sciuro-hypnum reflexum</i> (Starke) Ignatov & Huttunen	4	0.3	0.4	0.3		z	M, Se, R	1821
<i>Sciuro-hypnum starkei</i> (Brid.) Ignatov & Huttunen	10	0.5	0.3	0.1		z	Si, L, M, Se, R, F	1824
<i>Syntrichia ruralis ruralis</i> (Hedw.) F.Weber & D.Mohr	1	1.0	0.0	0.0		v	L	936
<i>Tetraphis pellucida</i> Hedw.	38	0.1	0.9	0.0		z	P, Si, L, T, M, Se, R, F	1575
<i>Thuidium assimile</i> (Mitt.) A. Jaeger	1	1.0	0.0	0.0		z-v	Si	1180
<i>Tortella tortuosa</i> (Hedw.) Limpr.	5	1.0	0.0	0.0		z-v	P, T, F	1647
<i>Tortula subulata</i> Hedw.	2	1.0	0.0	0.0		z	Si	1190
<i>Tritomaria exsecta</i> (Schmidel ex Schrad.) Loeske	6	0.4	0.5	0.1	NT	r	P, L, T, M, R	1572
<i>Tritomaria exsectiformis</i> (Breidl.) Loeske	2	0.7	0.3	0.0	NT	z	P, Si	1677

## Riassunto

Le conoscenze sulle briofite degli ambienti forestali sulle Alpi, e in particolare in Trentino-Alto Adige, sono ancora limitate. La maggior parte dei lavori sono floristici, e sono ancora assenti studi quantitativi che esaminano l'ambiente di pecceta. Gli obiettivi del presente lavoro sono: 1) accrescere le conoscenze sulla distribuzione delle briofite nelle peccete della provincia di Bolzano; 2) discutere l'ecologia e la distribuzione delle specie, o gruppi di specie, più significativi. In totale sono stati esaminati 48 plots lungo un gradiente altitudinale e in 8 aree della provincia. Le briofite sono state esaminate su tre diversi substrati, tronchi di abete rosso, legno morto e suolo. Il lavoro ha permesso di identificare 120 specie (91 di muschi e 29 di epatiche), 27 delle quali appartenenti alla Lista Rossa. Una interessante scoperta è stata la specie *Sciuro-hypnum curtum*, un muschio pleurocarpo segnalato per la prima volta in Italia. I principali fattori che determinano la distribuzione delle briofite in questi ambienti sono l'altitudine, la copertura forestale, la piovosità e la quantità di necromassa. Viene discusso il ruolo dell'ambiente e la responsabilità della gestione forestale per la conservazione della biodiversità.

## Acknowledgments

This work was funded by the Autonomous Province of Bolzano (Ripartizione Diritto allo studio, Università e Ricerca scientifica) in the framework of the project "Integrity assessment of South Tyrol forests by means of bryophytes distribution analysis". The Forest Planning Office of the Autonomous Province of Bolzano (project partner) is thanked for providing logistic and technical support. Special thanks to Petra Mair (Museum of Nature South Tyrol), Philipp Oberegger and Martin Stecher (Forest Planning Office of the Province) for field assistance. I'm in debt with Juri Nascimbene for the data collected in his project "Biodiversità, biomonitoraggio e conservazione dei licheni epifiti negli ambienti forestali della provincia di Bolzano", funded by the Autonomous Province of Bolzano. Lars Hedenäs (Swedish Museum of Natural History, Stockholm) and Jan Kučera (University of South Bohemia) are warmly thanked for revision of many critical species. Katarzyna Buczkowska (Adam Mickiewicz University, Poznań) provided useful indications concerning the genus *Calypogeia*.

## References

- A.A. V.V., 2010: Tipologie forestali dell'Alto Adige. Ripartizione per le Foreste vol. 2. Provincia Autonoma di Bolzano-Alto Adige.
- ALEFFI M. & CORTINI PEDROTTI C., 1996: Contributo alla flora briologica del gruppo del monte Bondone (Trentino-Alto Adige). *Informatore Botanico Italiano*, 28(2): 161-170.
- BUCZKOWSKA K., 2004: The genus *Calypogeia* Raddi (Jungermanniales, Hepaticae) in Poland, biometrical analysis of morphological and anatomical variation. *Nova Hedwigia*, 78: 121-146.
- CORTINI PEDROTTI C., 2001: Flora dei muschi d'Italia. Sphagnopsida, Andreaeopsida, Bryopsida (I parte). Antonio Delfino Editore, Roma.
- CORTINI PEDROTTI C., 2006: Flora dei muschi d'Italia. Bryopsida (II parte). Antonio Delfino Editore, Roma.
- CORTINI PEDROTTI C. & ALEFFI M., 2011: Lista Rossa delle Briofite del Trentino. *Studi Trentini di Scienze Naturali*, 88: 5-27.
- DALLA TORRE K.W. & SARNTHEIN L., 1904: Flora der gefürsteten Grafschaft Tirol, des Landes Vorarlberg und des Fürstentums Liechtenstein. Bd. 5: Die Moose (Bryophyta) von Tirol, Vorarlberg und Liechtenstein. Wagner, Innsbruck.
- DAMSHOLT K., 2002: Illustrated flora of Nordic liverworts and hornworts. Lund: Nordic Bryological Society.
- DULL R., 2006: Provisorischer Katalog der Leber- und Laubmoose Südtirols (Provinz Bozen). *Gredleriana*, 6: 69-114.
- GRYTNES J.A., HEEGAARD E. & IHLEN P.G., 2006: Species richness of vascular plants, bryophytes and lichens along an altitudinal gradient in western Norway. *Acta Oecologica*, 29: 241-246.
- HEDENÄS L. & BISANG I., 2004: Key to European Dicranum species. *Herzogia*, 17: 179-197.
- HUSCH B., BEERS, T.W. & KERSHAW, J.A., 2003: Forest mensuration. John Wiley & Sons, Inc.
- IGNATOV M. & MILYUTINA I.A., 2007: A revision of the genus *Sciurohypnum* (Brachytheciaceae, Bryophyta) in Russia. *Arctoa*, 16: 63-86.
- KIRÁLY I. & ÓDOR P., 2010: The effect of stand structure and tree species composition on epiphytic bryophytes in mixed deciduous-coniferous forests of Western Hungary. *Biological Conservation*, 143(9), 2063-2069.
- KOLARI P., PUMPANEN J., KULMALA L., ILVESNIEMI H., NIKINMAA E., GRÖNHOLM T. & HARI P., 2006: Forest floor vegetation plays an important role in photosynthetic production of boreal forests. *Forest Ecology and Management*, 221: 241-248.
- KORHONEN L., KORHONEN K.T., RAUTIAINEN M. & STENBERG P., 2006: Estimation of forest canopy cover: a comparison of field measurement techniques. *Silva Fennica*, 40(4): 577-588.
- JONSSON B.G., KRUYIS, N. & RANIS T., 2005: Ecology of species living on dead wood—lessons for dead wood management. *Silva Fennica*, 39(2): 289-309.
- LINDENMAYER D.B., MARGULES C.R. & BOTKIN D.B., 2000: Indicators of Biodiversity for Ecologically Sustainable Forest Management. *Conservation Biology*, 14: 941-950.
- MÁRIALIGETI S., NÉMETH B., TINYA F. & ÓDOR P., 2009: The effects of stand structure on ground-floor bryophyte assemblages in temperate mixed forests. *Biodiversity and conservation*, 18(8): 2223-2241.
- MILLS S.E. & MACDONALD S.E., 2005: Factors influencing bryophyte assemblage at different scales in the Western Canadian boreal forest. *Bryologist*, 108: 86-100.
- MOTTA R., BERRETTI R., LINGUA E. & PIUSSI P., 2006: Corse woody debris, forest structure and regeneration in the Valbona Forest Reserve, Paneveggio, Italian Alps. *Forest Ecology and Management*, 235:155-163.
- MÜLLER J. & BÜTLER R., 2010: A review of habitat thresholds for dead wood: a baseline for management recommendations in European forests. *European Journal of Forest Research*, 129: 981-992.
- NASCIMBENE J., 2014: Increasing the knowledge on the epiphytic lichens of South Tyrol: a contribution from a three-years project. *Gredleriana*, 14: 111- 126.
- NASCIMBENE J., NIMIS P.L. & DAINESI M., 2014: Epiphytic lichen conservation in the Italian Alps: the role of forest type. *Fungal Ecology*, 11: 164-172.
- NASCIMBENE J. & MARINI L., 2015: Epiphytic lichen diversity along elevational gradients: biological traits reveal a complex response to water and energy. *Journal of Biogeography*, DOI: 10.1111/jbi.12493
- NYHOLM E., 1986: Illustrated flora of Nordic mosses. Nordic Bryological Society
- OECHEL W.C. & VAN CLEVE K., 1986: The role of bryophytes in nutrient cycling in the Taiga. In: VAN CLEVE K., CHAPIN F.S. III, FLANAGAN P.W., VIERECK L.A. & DYRNESS C.T. (eds): *Forest ecosystems in the Alaskan taiga*. Springer Berlin Heidelberg, New York: 121-137
- PATON J.A., 1999. *Liverwort flora of the British Isles*. Harley Books.
- PROCTOR M.C.F., 2000: The bryophyte paradox: Tolerance of desiccation, evasion of drought. *Plant Ecology*, 151: 41-49.
- R DEVELOPMENT CORE TEAM, 2013: R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- ROS R.M., MAZIMPAKA V., ABOU-SALAMA U., ALEFFI M., BLOCKEEL T.L., BRUGUES M., CROS R.M., DIA M.G., DIRKSE G.M., DRAPER I., EL SAADAWI W., ERDAG A., GANEVA A., GABRIEL R., GONZALEZ-MANCEBO J.M., GRANGER C., HERRNSTADT I., HUGGONOT V., KHALIL K., KÜRSCHNER H., LOSADA-LIMA A., LUIS L., MIFSUD S., PRIVITERA M., REFAI M.S., SABOVLEVIC M., SERGIO C., SHABBARA H., SIM-SIM M., TACCHI R., VANDERPOORTEN A. & O. WERNER, 2013: Mosses of the Mediterranean, an annotated checklist. *Cryptogamie, Bryologie*, 34(2): 99-283.
- ROS R.M., MAZIMPAKA V., ABOU-SALAMA U., ALEFFI M., BLOCKEEL T.L., BRUGUÉS M., CANO M.J., CROS R.M., DIA M.G., DIRKSE G.M., EL SAADAWI W., ERDAG A., GANEVA A., GONZÁLES-MACEBO J.M., HERRNSTADT I., KHALIL K., KÜRSCHNER H., LANFRANCO E., LOSADA-LIMA A., REFAI M.S., RODRÍGUEZNUÑEZ S., SABOVLEVIC M., SERGIO C., SHABBARA H., SIMSIM M. & SÖDERSTRÖM L., 2007: Hepatics and Anthocerotae of the Mediterranean, an annotated checklist. *Cryptogamie, Bryologie*, 28: 351-437.
- SMITH A.J.E., 2004: *The moss flora of Britain and Ireland*. Cambridge University Press.

- SÖDERSTRÖM L., 1988: Sequence of bryophytes and lichens in relation to substrate variables of decaying coniferous wood in northern Sweden. *Nordic Journal of Botany* 8: 89–97.
- SPITALE D., 2015a: *Sciuro-hypnum curtum*. *Journal of Bryology*, 37 (4): in press.
- SPITALE D. 2015b: The interaction between elevational gradient and substrata reveals how bryophytes respond to the climate. *Journal of Vegetation Science*, submitted.
- SPITALE D. & MAIR P., 2015: Predicting the distribution of a rare species of moss: the case of *Buxbaumia viridis* (Bryopsida, Buxbaumiaceae). <http://dx.doi.org/10.1080/11263504.2015.1056858>
- SPITALE D., MAIR P. & TRATTER W., 2015: *Buxbaumia viridis* (Bryopsida, Buxbaumiaceae) in Alto Adige: un buon bioindicatore della quantità di necromassa in pecceta. *Gredleriana*, 15: xxx-yyy.
- SVEINBJORNSSON B. & OECHEL W.C., 1992: Controls on growth and productivity of bryophytes: environmental limitations under current and anticipated conditions. In: BATES J.W. & FARMER A.W. (eds): *Bryophytes and lichens in a changing environment*. Clarendon Press, Oxford: 77-102.
- TIERNEY G.L., FABER-LANGENDOEN D., MITCHELL B.R., SHRIVER G. & GIBBS J., 2009: Monitoring and evaluating the ecological integrity of forest ecosystems. *Frontiers in Ecology and the Environment*, 7: 308–316.
- TINYA F., MÁRIALIGETI S., KIRÁLY I., NÉMETH B. & ÓDOR P., 2009: The effect of light conditions on herbs, bryophytes and seedlings of temperate mixed forests in Orség, Western Hungary. *Plant Ecology*, 204: 69–81.